April 10, 2018

Honorable Chairman Jeb Bradley Senate Health and Human Services Committee Room 101 Legislative Office Building 33 N. State Street Concord, NH 03301

Dear Chairman Bradley and Members of the Committee,

Thank you for the opportunity to testify today. My name is Kenneth Norton and I serve as Executive Director of NAMI NH, the National Alliance on Mental Illness. I also have a family member with a serious mental illness and co-occurring substance use disorder. On behalf of NAMI NH, I am here to speak in support of HB 587. I am also here today to speak as a Licensed Independent Clinical Social Worker (LICSW) in the state of NH and as member of the National Association of Social Workers (NASW). NASW has opposed conversion therapy since 1992 as having no basis in science and being in direct conflict with the NASW Code of Ethics.

One of the maxims that all health professionals should aspire to is to *do no harm*. The potential for harm caused by conversion therapy rises beyond shame and trauma; which can have lifelong impacts, and has the potential to be life threatening. As the chart below indicates, suicide is the

second leading cause of death for ages 10-24 in NH (as well as nationally). According to the US Center for Disease Control, LGBT youth represent a significant high risk sub-group as evidenced by their rate of suicide attempts which is 4 times greater for LGB youth and 2 times greater for questioning youth, than that of straight youth. Additionally, suicide attempts by LGB youth and questioning youth are 4 to 6 times more likely to result in injury, poisoning, or overdose that requires treatment from a doctor or nurse, compared to their straight peers. Data has been

NH Top Ten Leading Causes of Death 2011-2015

		Age Groups									
Rank	≤1	1-4	5-9	10-14	15-24	<u>25-34</u>	<u>35-44</u>	<u>45-54</u>	<u>55-64</u>	<u>65+</u>	All Age
1	Conganital Anomalies 44	Unintentional Injury 10	Malignant Neoplasms	Malignant Neoplasms 11	Unintentional Injury 252	Unintentional Injury 458	Unintentional Injury 381	Malignant Neoplasms 993	Malignant Neoplasms 2,572	Heart Disease 10,179	Malignar Neoplasm 13,465
2	Short Gestation 41	Congenital Anomalies	Unintentional Injury 	Suicide 	Suicide 109	Suicide 149	Malignant Neoplasms 207	Heart Disease 870	Heart Disease 1,195	Malignant Neoplasms 9,561	Heart Disease 12,126
3	Maternal Pregnancy Comp. 29	Heari Disease	Homicide 	Unintentional Injury 	Mailignant Neoplasms 20	Matignant Neoplasma 7d	Sulcide 166	Unintentional Injury 453	Unintentional Injury 355	Chronic Low. Hespiratory Disease 2,936	Chronic Lo Respirato Disease 3,377
4	Placenta Cord Membranea 17	Homicide 	Carabro- vasqular	Benign Neoplasms	Meart Disease 14	Heart Disease 38	Heart Disease 119	Suicide 281	Chronic Low. Respiratory Disease 329	Carabro- vascular 2,082	Unintentio Injury 3,276
5	3105 11	Influenza & Pneumonia —	Congenital Anomalies —	Congenital Anomalies	Homicide 14	Congenital Anomalies 14	Liver Disease 37	Liver Disease 188	Liver Disease 250	Alahaimer's Disease 1,924	Cerebro vascular 2,319
6	Circulatory System Disease 10	Matignant Neoplasms	Heart Disease 	Heart Disease	Cerebro- vascular	Homicide 12	Diabeles Mellitus 23	Diabetes Mellitus 100	Diabetes Mellitus 248	Unintentional Injury 1,349	Alaheima Diseasa 1,966
7	Respiratory Distress	Benign Neoplasms 	Suicide —	Homicide 	Chronic Low. Respiratory Disease	Diabates Mellitus	Homicide 17	Chronic Low. Respiratory Disease 85	Suicide 263	Diabetes Mellitus 1,083	Diabete: Mellitus 1,473
8	Necrotizing Enteropolitis				Benign Neoplasms 	Chronic Low Respiratory Disease 	Carabro- yasoular 13	Carebro- vascular 64	Carebro- vascular 144	Influenza & Pnaumonia 1,022	Influenza & Pneumo 1,119
9	Neonatal Hamorhaga 				Congenital Anomalies	Liver Disease	Chronio Low. Respiratory Disease 13	Viral Hapatitis 27	Septicemia 73	Nephritis 782	Suicide 1,060
10	Unintentional Injury 				Diabetes Mellitus	Complicated Pregnancy	Two Tied	Influenza & Pneumonia 24	Viral Hepatitis 69	Parkinson's Disease 582	Nephriti 869

Terms for Causes of Death

Printer-Friendly Version

much more difficult to obtain specific to transgender youth, but all current evidence points to their being at much higher risk.

Find Help, Find Hope

However, more closely related to this bill, and the potential for harm of conversion therapy, is that LGB youth who come from highly rejecting families are 8.4 times as likely to have attempted suicide as LGB peers who reported no or low levels of family rejection.

In addition, the Journal of the American Medical Association (JAMA) released a study in 2017 showing decreased suicide rates both for GLB teens and for the teen population overall in states that have passed gay marriage.

The practice of conversion therapy in any form is a danger to the mental health and well-being of youth and young adults. Please vote HB 587 as ought to pass. Thank you for your consideration.

Respectfully,

Kenneth Norton, LICSW

Executive Director